

UTILITY PATENT APPLICATION
UNDER 37 CFR 1.53 (b)HONORABLE COMMISSIONER OF
PATENTS AND TRADEMARKS
Washington D.C. 20231Case Docket No. 1459-005B

Sir:

Transmitted herewith for filing is the patent application of

INVENTOR: Yoke Min Sin, Toong Jin Lam, Zhiyuan Gong

FOR: A Recombinant Vaccine Against Fish Infectious Diseases

Enclosed are:

- ☒ 25 pages of specification, claims, abstract
- ☒ Declaration & Power of Attorney.
- ☒ Priority Claimed
- ☐ Certified copy of _____
- ☐ _____ sheets of formal drawing.
- ☒ An assignment of the invention to National University of Singapore _____ and the assignment recordation fee.
- ☒ Return Receipt Postcard
- ☒ A verified statement to establish small entity status under 37 CFR 1.9 and 37 CFR 1.27.
- ☐ Information Disclosure Statement, Form PTO-1449
- ☐ Copies of IDS citations
- ☒ Machine Readable - Sequence listing disk

The filing fee has been calculated as shown below:

(1) FOR	(2) NO. FILED	(3) NO. EXTRA	(4) RATE	(5) AMOUNT
TOTAL CLAIMS	23	-20	3	x \$11.00=\$ 33.00
INDEPENDENT CLAIMS		-3	4	x \$41.00=\$161.00
MULTIPLE DEPENDENT CLAIM(S) (If applicable)				+ \$130.00 = 00.00
			BASIC FEE	\$ 395.00
Total of above calculations				= \$589.00
<input checked="" type="checkbox"/> Assignment & Recording Fee				\$40.00
			TOTAL FEE	\$629.00

11/20/98
14595 U.S. PTO11/20/98
14595 U.S. PTO
09/19/98

09/19/98 14595 U.S. PTO

	1970	1971	1972	1973	1974	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024	2025	2026	2027	2028	2029	2030	2031	2032	2033	2034	2035	2036	2037	2038	2039	2040	2041	2042	2043	2044	2045	2046	2047	2048	2049	2050	2051	2052	2053	2054	2055	2056	2057	2058	2059	2060	2061	2062	2063	2064	2065	2066	2067	2068	2069	2070	2071	2072	2073	2074	2075	2076	2077	2078	2079	2080	2081	2082	2083	2084	2085	2086	2087	2088	2089	2090	2091	2092	2093	2094	2095	2096	2097	2098	2099	2100	2101	2102	2103	2104	2105	2106	2107	2108	2109	2110	2111	2112	2113	2114	2115	2116	2117	2118	2119	2120	2121	2122	2123	2124	2125	2126	2127	2128	2129	2130	2131	2132	2133	2134	2135	2136	2137	2138	2139	2140	2141	2142	2143	2144	2145	2146	2147	2148	2149	2150	2151	2152	2153	2154	2155	2156	2157	2158	2159	2160	2161	2162	2163	2164	2165	2166	2167	2168	2169	2170	2171	2172	2173	2174	2175	2176	2177	2178	2179	2180	2181	2182	2183	2184	2185	2186	2187	2188	2189	2190	2191	2192	2193	2194	2195	2196	2197	2198	2199	2200	2201	2202	2203	2204	2205	2206	2207	2208	2209	2210	2211	2212	2213	2214	2215	2216	2217	2218	2219	2220	2221	2222	2223	2224	2225	2226	2227	2228	2229	2230	2231	2232	2233	2234	2235	2236	2237	2238	2239	2240	2241	2242	2243	2244	2245	2246	2247	2248	2249	2250	2251	2252	2253	2254	2255	2256	2257	2258	2259	2260	2261	2262	2263	2264	2265	2266	2267	2268	2269	2270	2271	2272	2273	2274	2275	2276	2277	2278	2279	2280	2281	2282	2283	2284	2285	2286	2287	2288	2289	2290	2291	2292	2293	2294	2295	2296	2297	2298	2299	2300	2301	2302	2303	2304	2305	2306	2307	2308	2309	2310	2311	2312	2313	2314	2315	2316	2317	2318	2319	2320	2321	2322	2323	2324	2325	2326	2327	2328	2329	2330	2331	2332	2333	2334	2335	2336	2337	2338	2339	2340	2341	2342	2343	2344	2345	2346	2347	2348	2349	2350	2351	2352	2353	2354	2355	2356	2357	2358	2359	2360	2361	2362	2363	2364	2365	2366	2367	2368	2369	2370	2371	2372	2373	2374	2375	2376	2377	2378	2379	2380	2381	2382	2383	2384	2385	2386	2387	2388	2389	2390	2391	2392	2393	2394	2395	2396	2397	2398	2399	2400	2401	2402	2403	2404	2405	2406	2407	2408	2409	2410	2411	2412	2413	2414	2415	2416	2417	2418	2419	2420	2421	2422	2
--	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	---

[X] Any additional filing fees required under 37 CFR 1.16.

[X] Any patent application processing fees under 37 CFR 1.17.

LOWE HAUPTMAN GOPSTEIN
GILMAN & BERNER, LLP

GILMAN & BERNER, LLP

 Michael G. Gilman

2304 Eisenhower Avenue, Suite 200
Alexandria, Virginia 22314
(703) 684-1111 MGG:jlg
FAX: (703) 518-5499
November 20, 1998

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re Application of :
Yoke Min Sin, Toong Jin Lam, Zhiyuan Gong :
Serial No. :
Filed: November 20, 1998 :
For: A Recombinant Vaccine Against Fish Infectious Diseases :

TRANSMITTAL OF VERIFIED STATEMENT
CLAIMING SMALL ENTITY STATUS

Honorable Assistant Commissioner
for Patents
Washington, D.C. 20231


Sir:

Transmitted herewith for filing in the above-referenced
application is(are) the following:

VERIFIED STATEMENT CLAIMING SMALL ENTITY STATUS -

Respectfully submitted,

LOWE HAUPTMAN GOPSTEIN
GILMAN & BERNER, LLP


Michael G. Gilman
Registration No. 19,114

2034 Eisenhower Avenue, Suite 200
Alexandria, Virginia 22314
(703) 684-1111 MGG:jljg
FAX: (703) 518-5499
Date: November 20, 1998

RECEIVED: 11/24/98

Applicant or Patentee: Yoke Min SIN, et al.Attorney's Docket No.: 1459-005B

Serial or Patent No.: _____

Filed or Issued: _____

For: A RECOMBINANT VACCINE AGAINST FISH INFECTIOUS DISEASES

**VERIFIED STATEMENT (DECLARATION) CLAIMING SMALL ENTITY STATUS
(37 CFR 1.9(f) and 1.27(d)) - NONPROFIT ORGANIZATION**

I hereby declare that I am an official empowered to act on behalf of the nonprofit organization identified below:

NAME OF ORGANIZATION: NATIONAL UNIVERSITY OF SINGAPOREADDRESS OF ORGANIZATION: 10 Kent Ridge Crescent, Singapore 119260

TYPE OF ORGANIZATION:

- ☒ UNIVERSITY OR OTHER INSTITUTION OF HIGHER EDUCATION
☐ TAX EXEMPT UNDER INTERNAL REVENUE SERVICE CODE (26 USC 501(a) AND 501(c)(3))
☐ NONPROFIT SCIENTIFIC OR EDUCATIONAL UNDER STATUTE OF STATE OF THE UNITED STATES OF AMERICA
(Name of State: _____)
(Citation of Statute: _____)
☐ WOULD QUALIFY AS TAX EXEMPT UNDER INTERNAL REVENUE SERVICE CODE (26 USC 501(a) and 501(c)(3)) IF LOCATED IN THE UNITED STATES OF AMERICA
☒ WOULD QUALIFY AS NONPROFIT SCIENTIFIC OR EDUCATIONAL UNDER STATUTE OF STATE OF THE UNITED STATES OF AMERICA IF LOCATED IN THE UNITED STATES OF AMERICA
(Name of State: _____)
(Citation of Statute: _____)

I hereby declare that the nonprofit organization identified above qualifies as a nonprofit organization as defined in 37 CFR 1.9(c) for purposes of paying reduced fees under section 41(a) and (b) of Title 35, United States Code with regard to the invention entitled A RECOMBINANT VACCINE AGAINST FISH INFECTIOUS DISEASES by inventor(s) Yoke Min SIN, Toong Jin LAM and Zhiyuan GONG described in

- ☐ the specification filed herewith
☐ application Serial No., filed.
☐ patent no., issued.

I hereby declare that rights under contract or law have been conveyed to and remain with the nonprofit organization with regard to the above identified invention.

If the rights held by the nonprofit organization are not exclusive, each individual, concern or organization having rights to the invention is listed below* and no rights to the invention are held by any person, other than the inventor, who could not qualify as a small business concern under 37 CFR 1.9(d) or by any concern which would not qualify as a small business concern under 37 CFR 1.9(d) or by a nonprofit organization under 37 CFR 1.9(e). *NOTE: Separate verified statements are required from each named person, concern or organization having rights to the invention averring to their status as small entities. (37 CFR 1.27)

NAME National University of Singapore
ADDRESS 10 Kent Ridge Crescent, Singapore 119260☐ INDIVIDUAL ☐ SMALL BUSINESS CONCERN ☒ NONPROFIT ORGANIZATION

I acknowledge the duty to file, in this application or patent, notification of any change in status resulting in loss of entitlement to small entity status prior to paying, or at the time of paying, the earliest of the issue fee or any maintenance fee due after the date on which status as a small entity is no longer appropriate. (37 CFR 1.28(b))

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under section 1001 of Title 18 of the United States Code, and that such willful false statements may jeopardize the validity of the application, any patent issuing thereon, or any patent to which this verified statement is directed.

NAME OF PERSON SIGNING: CHOU Siow Kang (Dr)TITLE IN ORGANIZATION: Director, Industry & Technology Relations Office (INTRO)ADDRESS OF PERSON SIGNING: National University of Singapore,
10 Kent Ridge Crescent, Singapore 119260

SIGNATURE: _____

DATE: 1 Oct. 1998

A RECOMBINANT VACCINE AGAINST INFECTIOUS DISEASE IN FISH

TECHNICAL FIELD

This invention is directed to a recombinant vaccine for immunizing fish against white spot disease (also known as ich).

BACKGROUND OF THE INVENTION

White spot disease is a severe infectious disease that adversely affects warm water fish. . It is caused by a ciliated ectoparasitic protozoan, *Ichthyophthirius multifiliis* that causes high morbidity and mortality in many freshwater fish. In the past, when fish show clinical signs of infection by this disease, they have commonly been treated with chemicals (Cross, D.G., "A REVIEW OF METHODS TO CONTROL ICHTHYOPHTHIRIASIS", Prog. Fish-culture, 34, (1972) 165-169; Gratzek, J.B., "CONTROL AND THERAPY OF FISH DISEASES", Advances Vet. Sci. Comparat. Med., 27, (1983) 297-324; Post, G., "TEXTBOOK OF FISH HEALTH" (Revised and expanded edition), T.F.H. Publications Inc, USA, 1987). However, these chemicals usually leave residues in the fish tissues that can endanger not only the health of the fish, but they can also adversely affect humans who are the ultimate consumers of the infected fish.

On the other hand, protection of many fish species against Ichthyophthiriasis has been reported through the use of immunization with *I. Multifiliis*. Such reports of immunized fish include:

goldfish (Ling, K.H., Sin, Y.M. and Lam, T.J., "PROTECTION OF GOLDFISH AGAINST SOME COMMON ECTOPARASITIC PROTOZOANS USING *ICHTHYOPHTHIRIUS MULTIFILIIS* AND *TETRAHYMENA PYRIFORMIS* FOR VACCINE", Aquaculture, 116 (1993) 303-314;

mirror carp (Hines, R.S. and Spira, D.T., "ICHTHYOPHTHIRIASIS IN THE MIRROR CARP *CYPRINUS CARPIO* (L), IV. PHYSIOLOGICAL DISFUNCTION", J. Fish Biol., 6 (1974) 365-371;

channel catfish (Gover, B.A., Dawe, D.L. and Gratzek, B., "PROTECTION OF CHANNEL CATFISH, *ICTALURUS PUNCTATUS* RAFINESQUE, AGAINST *I. MULTIFILIIS* FOUQUET BY IMMUNIZATION", J.Fish Biol., 17 (1980) 311-316 ;

rainbow trout (Wolf, K. and Markiw, M.E., "ICHTHYOPHTHIRIASIS: IMMERSION IMMUNIZATION OF RAINBOW TROUT (*SALMO GAIRDNERI*) USING *TETRAHYMENA THERMOPHILA* AS A PROTECTIVE IMMUNOGEN", Can. J. Fish Aquat. Sci., 39 (1982) 1772-1725;

tilapia (Subasinghe, R.P. and Sommerville, C., "ACQUIRED IMMUNITY OF *OREOCHROMIS MOSSAMBISCUS* TO THE CILIATE ECTOPARASITE *ICHTHYOPHTHIRIUS MULTIFILIIS* (FOUQUET)" In : J.L Maclean, L.B. Dizon and L.V. Osillos (editors), The first Asian Fisheries Forum, Asean Fisheries Society, Manila, Philippines, 1987, 279-283); and

some other common ornamental fish (Ling K.H., Sin Y.M. and Lam T.J., "A NEW APPROACH TO CONTROLLING ICHTHYOPHTHIRIASIS IN A CLOSED CULTURE SYSTEM OF FRESHWATER ORNAMENTAL FISH", J. fish Dis., 14 (1991) 595-598).

Until now, no known methods have been reported as to how to *in vitro* grow the protozoan, *I. Multifiliis*, that has caused the problem. The vaccines (whole cells, cell fragments or extracts of the pathogen), that have been prepared and used by different laboratories, have only been obtained from an infected live fish since the offending protozoan is an obligate parasite. Preparation of such vaccines from live fish is very time-consuming, costly and impractical.

An antigenic relationship has been demonstrated between *I. Multifiliis* and *Tetrahymena pyriformis*, a free-living and cultivable non-pathogenic ciliated protozoan (Goven, B.A., Dawe, D.L. and Gratzek, J.B., "PROTECTION OF CHANNEL CATFISH, *ICTALURUS PUNCTATUS* RAFINESQUE, AGAINST *ICHTHYOPHTHIRIUS MULTIFILIIS* FOUQUET BY IMMUNIZATION", J. Fish Biol., 17 (1980) 311-316). These workers suggested that *T. pyriformis* could be used as a vaccine for inducing protection against Ichthyophthiriasis (Goven, B.A., Dawe, D.L. and Gratzek, J.B., "IN VITRO DEMONSTRATION OF SEROLOGICAL CROSS-REACTIVITY BETWEEN *ICHTHYOPHTHIRIUS MULTIFILIIS* FOUQUET AND *T. PYRIFORMIS* LWOFF", Dev. Comp. Immunol., 5 (1981) 283-289). A ciliary protein extracted from *T. pyriformis* was shown to be an effective vaccine against the disease caused by *I. Multifiliis* in freshwater fish and *Cryptocryon* in marine fish (U.S. Patent Number 4,309,416). The presence of an immobilization antigen (i-Ag) in *I. Multifiliis* was

reported by Dickerson, H.W., Clark, T.G. and Findly, R.C., "*ICHTHYOPHTHIRIUS MULTIFILIIS* : HAS MEMBRANE-ASSOCIATED IMMOBILIZATION ANTIGENS", J. Protozool., 36 (1989) 159-164) and by Lin, T.L. and Dickerson, H.W., "PURIFICATION AND PARTIAL CHARACTERIZATION OF IMMOBILIZATION ANTIGEN FROM *ICHTHYOPHTHIRIUS MULTIFILIIS*", J. Protozool., 39 (1992) 457-463).

A partial cDNA clone for 48 kDa i-Ag was isolated from a cDNA library and the amino acid sequence that was deduced from the cDNA clone contained three tandem repeats (Clark, T.G., McGraw, R.A. and Dickerson, H.W., "DEVELOPMENTAL EXPRESSION OF SURFACE ANTIGEN GENES IN THE PARASITIC CILIATE *ICHTHYOPHTHIRIUS MULTIFILIIS*", Proc. Natl. Acad. Sci. USA, 89 (1992) 6363-6367). In fact, the degree of immune protection was correlated with the levels of antibody titers against this immobilization antigen in the plasma and skin mucus of the fish that had been immunized against *I. Multifiliis* (Clark, T.G., Dickerson, H.W., and Findly, R.C., "IMMUNE RESPONSE OF CHANNEL CATFISH TO CILIARY ANTIGENS OF *ICHTHYOPHTHIRIUS MULTIFILIIS*", Devel. Comp. Immunol., 12 (1988) 581-594 and *T. pyriformis* (Ling, K.H., Sin, Y.M. and Lam T.J., "PROTECTION OF GOLDFISH AGAINST SOME COMMON ECTOPARASITIC PROTOZOANS USING *ICHTHYOPHTHIRIUS MULTIFILIIS* AND *TETRAHYMENA PYRIFORMIS* FOR VACCINATION", Aquaculture, 116 (1993) 303-314). Fish immunized with *I. Multifiliss* and *T. pyriformis* by immersion and intraperitoneal injection developed protective immunity not only against *I. Multifiliis*, but also against other ciliated ectoparasitic protozoans that are commonly found in the tropics (Ling, K.H., Sin, Y.M. and Lam T.J., "PROTECTION OF GOLDFISH AGAINST SOME COMMON ECTOPARASITIC

PROTOZOANS USING *ICHTHYOPHTHIRIUS MULTIFILIIS* AND *TETRAHYMENA PYRIFORMIS* FOR VACCINATION”, Aquaculture, 116 (1993) 303-314.

Since the chemicals that have been found to be effective against these adverse ciliated protozoans are not approved for use in food fish by the USDA, and many other countries in the world, immunization may be the only practical approach to controlling this white spot disease. This can be done by either direct immunization of the fish with vaccines, using whole cells or ciliary protein extracts from *I. multifiliis*, as discussed above or by passive protection using immobilization antibodies against the pathogen released by the immunized fish in a closed culture system (U.S. Patent Number 5,643,571).

OBJECTS AND BRIEF DESCRIPTION OF THE INVENTION

An important object of this invention is to provide an effective vaccine for immunization against infectious fish ectoparasitic protozoan, *Ichthyophthirius mutifiliis* (He, J., Yin, Z., Xu, G., Gong, Z., Lam, T.J. and Sin, Y.M., “PROTECTION OF GOLDFISH AGAINST *ICHTHYOPHTHIRIUS MULTIFILIIS* BY IMMUNIZATION WITH A RECOMBINANT VACCINE”, Aquaculture, 158(1997)1-10) that has been produced by recombinant techniques, since no vaccine has as yet been obtained, nor is it likely that one can be obtained, using *in vitro* systems.

Another object of this invention is to provide an effective vaccine for the immunization of fish, not only against *Ichthyophthirius multifiliis*, but also against other, taxonomically related, ciliated ectoparasitic protozoans, by recombinant techniques.

A further object of this invention is to provide a feasible method for commercial production of this vaccine.

Other and additional objects will become apparent from a consideration of this entire specification including the claims appended hereto.

In accordance with and fulfilling these objects, one aspect of this invention is a vaccine against ciliated protozoan infection that has been prepared by recombinant techniques. A gene fragment, containing a complete repeat I of *I. Multifiliis*, was synthesized and assembled based on the amino acid sequence of i-AgI. The recombinant i-AgI was expressed in bacteria, as a glutathione S-transferase (GST-iAgI) fusion protein. The fusion protein was purified from bacterial lysate and successfully used as a vaccine for immunization. The modified sequences according to this invention can be used to produce the recombinant fusion protein by any organism. These are exemplified by: yeast, baculovirus in insect cells, transgenic mammals, other vertebrate and invertebrate and transgenic plants using universal genetic codons. These universal codons are CAG and CAA, which are glutamine codons.

BRIEF DESCRIPTION OF THE DRAWING

Fig 1A is an illustration of the epitope of i-Ag included in a GST-iAgI recombinant protein;

Fig. 1B is an illustration of six synthetic oligonucleotides that were used to construct the i-AgI epitope region;

Fig. 1C is a diagram of the plasmid pGST-iAgI that was used for the production of GST-iAgI protein;

Figs. 2(A) is a plan view of a gel electrophoresis of bacterial expressions of GST-iAgI fusion protein; and

Fig 2(B) is a plan view of a Western Blot analysis.

DETAILED DESCRIPTION OF THE INVENTION

A 316 bp gene fragment containing an antigenic epitope of the 48K immobilization antigen of *Ichthyophthirius multifiliis* (iAgI) was assembled from six synthetic oligonucleotides and cloned into a bacterial expression vector pGEX2T. The gene construct was introduced into *Escherichia coli* and the glutathione S-transferase-iAgI (GST-iAgI) fusion protein was successfully expressed. Antisera against GST-iAgI fusion protein from catfish showed a positive

reaction with a tomite protein of about 48K, suggesting that the recombinant protein contains an antigenic epitope of i-AgI. Naive goldfish, that were immunized with purified GST-iAgI fusion protein, were challenged with a lethal dose of infectious tomites of *I. Multifiliis*. The results showed that the recombinant GST-iAgI fusion protein can be used as a vaccine against the infection of not only *I. Multifiliis* but also against diseases caused by other taxonomically similar pathogens, that is the ciliated ectoparasitic protozoans.

Referring now to Fig. 1(A), the amino acid sequence of the epitope of i-Ag is set forth on line I, the corresponding nucleotide sequences from the original i-Ag cDNA clone are set forth on line II, and the synthetic DNA fragment of this invention is set forth on line III. This Fig. aligns these three sequences. Two conserved amino acid substitutions between the original and the synthetic i-AgI DNAs have been indicated as A/S and V/G, where S and G have been derived from the synthetic gene. The amino acids from and including 23 to 101 have been marked off by square brackets. These are a repeat of i-Ag (see Clark et al., 1992, above). The glutamine codons TAA and TAG in the original sequence, which are the universal codons, are indicated by asterisks, and have been replaced with the universal glutamine codons CAG and CAA (underlined) in the synthetic gene. The restriction sites in the synthetic gene that are used for cloning have also been underlined. The changed nucleotides for the creation of the restriction enzyme sites, as well as for the replacement of the glutamine codons, have been shown in italics.

Referring now to Fig. 1(B), there is shown six (6) synthetic oligonucleotides that were used to construct the i-AgI epitope region. There are three (3) DNA segments, each of which is constructed with two (2) oligonucleotides that were derived from the coding strand (upper) and

the non-coding strand (lower). The 3' complimentary nucleotides (12bp) have been aligned, and the 5' restriction sites and changed glutamine codons have been underlined.

Referring now to Fig. 1(c), there is shown a diagram of the plasmid pGST-iAgI used for the production of GST-iAgI protein. The 316 bp synthetic DNA fragment was inserted into pGEX2T vector between BamHI and EcoRI sites as described herein.

Referring now to Fig. 2(A), there are shown bacterial expressions of GST-iAgI fusion protein. Lanes 1-4 show total lysate of bacteria transformed with pGEX2T (lanes 1 and 2), or with pGST-iAgI (lanes 3 and 4). Lanes 1 and 3 had no IPTG induction, and lanes 2 and 4 had IPTG induction. Lanes 5 shows purified GST-iAgI with thrombin digestion and lane 6 shows GST-iAgI without thrombin digestion. The GST-iAgI was cleaved into two (2) parts, as indicated. The SDS-PAGE was stained with coomassie brilliant blue. The position of the protein is indicated by the solid dot on the right of each lane. M was a prestained protein standard (Gilbco, Life Sciences) in KDa.

Referring now to Fig. 2(B), which is a Western Blot analysis, in lane 1 there is shown GST-iAgI fusion protein that has been partially digested with thrombin. The solid dots indicate intact fusion protein GST and i-AgI. In lane 2 there is shown the Tomite total protein. M indicates a high range of prestained protein standard (Gilbco, Life Sciences) in KDa.

MATERIALS AND METHODS

DESIGN, SYNTHESIS AND ASSEMBLY OF THE 'iAGI' GENE

The immobilization antigen of *I. Multifiliis* (i-AgI) consisted of three tandem repeats in its central region. Three DNA segments of similar length were designed to cover a complete repeat I of i-AgI. For each segment, two oligonucleotides, representing both coding and noncoding strands, were synthesized. To ensure efficient intermolecular annealing, the two oligonucleotides contained 12 complementary nucleotides at their 3' ends (Fig. 1B). Restriction sites were engineered at the 5' ends of each oligonucleotide to facilitate cloning. To generate double-stranded DNA from the oligonucleotides, 10 :1 each of the corresponding two oligonucleotides were mixed, at the same concentration (1:µg:1), with 5:1 of 10 X Klenow buffer (500mM Tris, pH7.2, 100mM MgSO₄, 1mM dithiothreitol) and 18:1 distilled water. The reaction mixture was heated and maintained at 80°C for 5 min and then allowed to cool to 37°C, followed by addition of 5 :1 of 2.5mM deoxyribonucleotides (dNTP) and 2 :1 of Klenow fragment (Promega) to initiate DNA synthesis at room temperature. After 40 minutes EDTA was added to a final concentration of 10 mM to stop the reaction, and this was followed by incubation at 75°C for 5 min to inactivate the Klenow enzyme. The double-stranded DNA segments were purified by phenol/chloroform extraction and ethanol precipitation, and dissolved in 50 :1 of distilled water. To assemble the three segments into a longer DNA fragment, half of double-stranded segments 1, 2 and 3 were double digested with BamHI-KpnI, KpnI-Sau3A1 and Sau3A1-EcoRI, respectively. The digested segments were recovered by gel purification, using low-melting-

temperature agarose, and dissolved in 25 :l of water. About 0.1 :g of digested segment 1 was ligated into 0.1 :g of pBluescript vector (Stratagene) that had been predigested with BamHI and KpnI. The ligation was performed overnight at 4°C in a 20 :l reaction mixture. Similarly, 0.2 :g each of segment 2 and segment 3 were ligated together into pBluescript vector (Stratagene) that had been predigested with KpnI and EcoRI. 10 :l of each ligation reaction mixture was transferred into 100 :l of *E. coli* competent cell (XL1-Blue strain, Stratagene) according to the standard protocol described by Sambrook, J., Fritsch E.F. and Maniatis, T. ("MOLECULAR CLONING, A LABORATORY MANUAL" 2nd edition, Cold Spring Harbor Laboratory Press, New York, 1989). The resulted plasmids were isolated with Qiagen Plasmid Isolation Kit (Qiagen) and sequenced from both ends for confirmation. The verified segment 1 and segments 2 and 3 were removed from pBluescript by digestion with BamHI-KpnI and KpnI-EcoRI respectively. 0.1 :g of verified segment 1 and segments 2 and 3 each were ligated into 0.1 :g of pGEX2T (Pharmacia) vector cut with BamHI and EcoRI. The volume of 10 :l ligation reaction was transformed into *E. coli* (XL1-Blue strain, Stratagene), and the resulting construct was called pGST-iAgI. Its sequence was confirmed again by sequencing with the primer 5'-TAGCATGGCCTTTGCAG-3', upstream of the cloning site of pGEX2T vector.

EXPRESSION AND PURIFICATION OF GST FUSION PROTEIN

An overnight culture of *E. coli* harboring pGST-iAgI was diluted to 1:20 in fresh LB medium containing 50 µg/ml ampicillin and grown at 37°C with vigorous shaking. Isopropyl 1-thio-β-D-galactoside (IPTG) was added to a final concentration of 0.1 mM when the OD₆₀₀

reached about 0.5. Bacteria were harvested, by centrifugation at 2,000 G at 4°C for 15 min, 3 hours after the addition of IPTG. The bacteria were resuspended in 300 µl of phosphate-buffered saline (PBS), boiled in SDS-sample buffer for 5 min and analyzed by electrophoresis (according to Laemmli, U.K., "CLEAVAGE OF STRUCTURE PROTEIN DURING THE ASSEMBLY OF THE HEAD OF BACTERIOPHAGE T4" Nature, 227(1970) 680-685)) using an SDS-polyacrylamide gel.

The GST-iAgI fusion protein was purified using Glutathione Sepharose 4B Beads (Pharmacia) as previously described (Gong, Z. and Hew, C.L., "ZINC AND DNA BINDING PROPERTIES OF A NOVEL LIM HOMEODOMAIN PROTEIN ISL-2", Biochemistry, 33 (1995)15149-15158). Shortly after IPTG induction, 1 liter of bacterial culture was harvested by centrifugation and resuspended in 10 ml of lysis buffer (50 mM Tris, pH8.0, 0.1M NaCl, 1mM EDTA), to which lysozyme had been added to a final concentration of 5 mg/ml. After incubation at room temperature for 5 min, bacteria, that were lysed with 1% Triton-100. MgCl₂ and DNase I, were then added, in amounts sufficient to produce final concentrations of 5mM and 10µg/ml respectively, to digest viscous DNA. The lysate was cleared by centrifugation. The supernatant was incubated with 1 ml of a 50% slurry of Glutathione Sepharose 4B Beads (Pharmacia) at room temperature for 1 hour with gentle agitation. The beads were then washed three times in 10X of bed volume of PBS. The GST-iAgI fusion protein was eluted from beads with 15 mM reduced glutathione (Sigma). The concentration of purified fusion protein was measured using Bio-Rad Protein Assay (Bio-Rad). Thrombin (Sigma) digestion of purified pGST-iAgI fusion protein was carried out according to the method described by Smith, D.B. and Johnson, K.S.

“SINGLE-STEP PURIFICATION OF POLYPEPTIDES IN *ESCHERICHIA COLI* AS FUSION GLUTATHIONE S-TRANSFERASE” Gene, 67 (1988)31-40.

ANALYSIS OF ANTIGENICTY OF GST-IAGI FUSION PROTEIN BY WESTERN BLOT.

Catfish, of 20-25 cm in length, were immunized by intraperitoneal injection of 200 µg of GST-iAgI fusion protein mixed with Freund's complete adjuvant (FCA). A similar dose of the fusion protein, without FCA, was repeated two weeks later. Pre-immunization sera were collected from each fish prior to immunization. Post immunization sera were collected one week after the second injection. Infectious tomites were collected according to the method as previously described (Ling, K.H, Sin, Y.M. and Lam, T.J., “PROTECTION OF GOLDFISH AGAINST SOME ECTOPARASITIC PROTOZOANS USING *ICHTHYOPHTHIRIUS MULTIFILIIS* AND *TETRAHYMENA PYRIFORMIS* FOR VACCINATION”, Aquaculture, 116 (1993)303-314).

The tomites and thrombin digested GST-iAgI fusion protein were boiled in SDS sample buffer for 5 min, and analyzed by electrophoresis using an SDS-polyacrylamide gel. The resolved protein was electroblotted onto a 0.2 µm nitrocellulose filter. For immunodetection, the filter was blocked in 0.5% bovine serum albumin (BSA) in TTBS (0.05% Tween- 20, 10 mM Tris-HCl, 150 mM NaCl, pH7.5) at 4°C for 16 hours, followed by incubating with 1:50 primary antibody (catfish antisera against GST-iAgI fusion protein) overnight at 4°C. After three washes with TTBS, the filter was incubated with 1:100 of rabbit anti-catfish serum for 1 hour and then with 1:3,000 dilution of goat anti-rabbit IgG alkaline phosphatase conjugates for 1 hour (Bio-Rad).

The filter was washed with TTBS and incubated with substrate NBT (nitroblue tetrazolium) and BCIP(5-bromo-4-chloro-3-indoyl phosphate) for color development at room temperature.

CHALLENGE TEST WITH INFECTIOUS TOMITES

Goldfish, *Carassius auratus* (L.), weighing 5-6 gms, were used throughout the study. All fish were bred from healthy broodstock and were not previously infected by *I. Multifiliis* nor by any other obvious parasites. The fish were maintained in the laboratory as described previously (Ling, K.H., Sin, Y.M. and Lam, T.J., "A NEW APPROACH TO CONTROLLING ICHTHYOPHTHIRIASIS IN A CLOSED CULTURE SYSTEM OF FRESHWATER ORNAMENTAL FISH", J. Fish Dis., 14 (1991) 595-598).

Each fish was immunized by intraperitoneal injection of 15 µg GST-iAgI fusion protein with FCS and boosted, by the injection of a similar dose of GST-iAgI fusion protein without the adjuvant, two weeks later. The control group was similarly injected with PBS and FCA mixture at the first injection, and then boosted with PBS alone. One week after the last injection, the control and immunized fish were challenged by direct exposure to infectious tomites using the method of Ling, K.H., Sin, Y.M. and Lam, T.J., "A NEW APPROACH TO CONTROLLING ICHTHYOPHTHIRIASIS IN A CLOSED CULTURE SYSTEM OF FRESHWATER ORNAMENTAL FISH", J. Fish Dis., 14(1991) 595-598. Each fish of all groups was exposed to 5,000 tomites. The fish were kept in aerated water at room temperature for four weeks under observation for infection and death. The degree of infection was separated and recorded in three (3) categories: none (free of parasites), mild, and heavy infection. Fish infected with less than

100 parasites per fish, and had parasites only on their fins were considered to have a mild infection. Those infected with more than 100 parasites, and not only on their fins but also on their body surface, were considered to have a heavy infection. The significance of the survival rate of fish between the immune and the control group after challenge was analysed by Chi-square test.

SPECIFIC EXAMPLES AND RESULTS

Example 1.

The amino acid sequence of i-AgI (Clark, T.G., McGraw, R.A. and Dickerson, H.W., "DEVELOPMENTAL EXPRESSION OF SURFACE ANTIGEN GENES IN THE PARASITIC CILIATE *ICHTHYOPHTHIRIUS MULTIFILIIS*", Proc. Natl. Acad. Sci. USA, 89 (1992) 6363-6367) was analyzed by a computer program to determine if it had a potential antigenic epitope, Predict 7 V1.01© (Carmenes, R.S., Freije, J.P., Molina, M.M. and Martin, J.M., "PREDICT 7, A PROGRAM FOR PROTEIN STRUCTURE PREDICTION", Biochem. Biophys. Res. Commun., 159 (1989) 687-693). It was predicted that each of the three repeats could be a strong antigenic epitope. The repeat I (Fig.1A) was selected for expression of the recombinant protein for immunization.

Three pairs of oligonucleotides were designed and synthesized to cover the repeat I of i-AgI (see Fig. 1). For the expression of i-AgI recombinant protein, the assembled gene fragment was inserted into the bacterial expression vector pGEX2T and the resulting plasmid was named pGST-iAgI (see Fig. 1). As shown in Fig.2(a), the expression of GST-iAgI fusion protein can be

induced after addition of IPTG (lane 4). The GST-iAgI fusion protein can be purified by glutathione Sepharose 4B bead (lane 6), and the purified GST-iAgI fusion protein can be cleaved at the fusion site with thrombin (lane 5). The yield of GST-ich fusion protein, purified from 1liter bacterial culture, was 2.7-3.0 mg.

Example 2.

To test whether the recombinant protein contained the antigenic epitope for i-AgI, antisera against GST-iAgI, fusion protein was raised in catfish and tested for immunological reaction with the total tomities' proteins. As shown in Fig. 2(b), the catfish antisera reacted strongly to the intact fusion protein, as well as to the GST and the iAgI portions after thrombin digestion (lane 1). It is interesting to note that the catfish antisera also reacted with a tomite's protein of a molecular weight of about 48K (lane 2). This is consistent with the predicted molecular weight of i-AgI. All the sera obtained from fish prior to immunization were negative (data not shown).

Example 3.

To test the effectiveness of the GST-iAgI as a vaccine, naive goldfish were immunized with the fusion protein and then challenged with infectious tomities of *I. Multifiliis*. Clinical signs of the disease and mortalities of the fish were recorded from day 4 onwards after the exposure to these tomities. The result are summarized in Table 1. 22.7% of the immunized fish showed no clinical sign of the disease, while 50% of the immunized fish exhibited heavy

infection, and 27.3 % showed a mild infection. In contrast, all the control fish were infected: 75% of non-immunized fish were heavily infected, and the other 25% were mildly infected. The average survival rate of the immunized fish was 95% as compared to 55% for the control. The difference between the survival rate of the control and of the immunized group is significant, $p < 0.001$ as determined by Chi-square-sequare test.

Table 1: Infectivity and average survival rate of control and immunized fish after being exposed to infective tomites of *I. Multifiliis*.

Group	Total fish *	<u>Degree of infectivity (%)**</u>			Survival (%)***
		heavy	mild	none	
Control	20	75	25	0	55%
Immune	22	50	27.2	22.8	95%

*Two duplicate groups of 10 to 12 fish each.

**Heavy : the number of parasites found on the fish body and fins was more than 100 up to 500-1,000; Mild: the number of parasites found only on the fins was less than 100; None: the fish were free of parasites

*** significantly different at $p < 0.001$ level, between control and immunized fish by the Chi-square test

FURTHER APPLICATION OF THE RECOMBINANT VACCINE

The effectiveness of this recombinant vaccine in immune protection against *Ichthyophthirius multifiliis* has been demonstrated in fish (He, J., Yin, Z, Xu, G., Gong Z., Lam T.J. and Sin Y.M., "PROTECTION OF GOLDFISH AGAINST *ICHTHYOPHTHIRIUS MULTIFILIIS* BY IMMUNIZATION WITH A RECOMBINANT VACCINE", Aquaculture, 158 (1997)1-10). The antigenicity of the vaccine was also shown using antisera against GST fusion protein from catfish that reacted strongly to the protein of *I. Multifiliis* of a similar size (48 KDa) to that of the immobilization antigen of *I. Multifiliis* (i-AgI) as reported by Clark, T.G., McGraw, R.A. and Dickerson, H.W., "DEVELOPMENTAL EXPRESSION OF SURFACE ANTIGEN GENES IN THE PARASITIC CILIATE *ICHTHYOPHTHIRIUS MULTIFILIIS* ", Proc. Natl. Acad. Sci. USA, 89 (1992) 6363-6367). In fact, fish immunized with the immobilization protein of *I. Multifiliis* and *T. pyriformis* showed protection against the disease caused by *I. Multifiliis* in freshwater fish (Goven , B.A., Dawe, D.L. and Gratzek, J.B., "PROTECTION OF CHANNEL CATFISH, *ICTALURUS PUNCTATUS* RAFINESQUE, AGAINST *ICHTHYOPHTHIRIUS MULTIFILIIS* FOUQUET BY IMMUNIZATION", J. Fish Biol., 17 (1980) 311-316; Goven,B.A., Dawe, D.L and Gratzek, J.B., " *IN VITRO* DEMONSTRATION OF SEROLOGICAL CROSS-REACTIVITY BETWEEN *ICHTHYOPHTHIRIUS MULTIFILIIS* FOUQUET AND *T. PYRIFORMIS* LWOFF", Dev. Comp. Immunol.,5 (1981) 283-289) and *Cryptocryon* in marine fish (U.S. Patent Number 4,309,416). The immune protection was correlated with the levels of immobilization antibody in the plasma and skin mucus of the freshwater fish immunized against *I. Multifiliis* or *T. pyriformis* (Clark , T.G., Dickerson, H.W., and Findly, R.C., "IMMUNE RESPONSE OF CHANNEL CATFISH TO CILIARY ANTIGENS OF *ICHTHYOPHTHIRIUS MULTIFILIIS*", Devel. Comp. Immunol.,12 (1988)581-594; Ling, K.H., Sin, Y.M. and Lam T.J., "PROTECTION OF

GOLDFISH AGAINST SOME COMMON ECTOPARASITIC PROTOZOANS USING *ICHTHYOPHTHIRIUS MULTIFILIIS* AND *TETRAHYMENA PYRIFORMIS* FOR VACCINATION", Aquaculture, 116 (1993) 303-314). An immobilization antigen extracted from *T. pyriformis* was also shown to be an effective vaccine against *I. Multifiliis* and *Cryptocryon* in marine fish (U.S, Patent Number 4,309,416). Since fish immunized with either *I. Multifiliis* or *T. pyriformis* developed protective immunity not only against *I. Multifiliis* but also against other ciliated ectoparasitic protozoans that are commonly found in the tropics, therefore, the recombinant vaccine made for the immobilization antigen of *I. Multifiliis* should also exert the same immune protection not only against *I. Multifiliis* but also against other taxonomically related ciliated ectoparasitic protozoans in the freshwater and the marine fish.

For immunization by injection, the fusion protein was dissolved in buffer and mixed with Freund's complete adjuvant (FCA). Any substance or medium that does not chemically destroy or interfere with the immunogenicity of the fusion protein can be incorporated into the formulation of the vaccine. These include, for example, buffers, immunostimulants and carriers. The vaccine can be introduced to fish by intraperitoneal injection, incorporated into fish diet, and by immersion or by spraying of the fish. For injection of a 5-6 gm fish, at least about 15 :g of the fusion protein is required. That is, a dose rate of at least about 3 :g fusion protein per gram of fish is appropriate

SEQ. ID NO 1

G A/S A Q* G E A N G N Q* P F A A N N A A R G I[C V P C Q I N R V G S V/ G T N A G D
L A T L A T Q* C S T Q* C P T G T A L D D G V T D V F D R S A A Q* C V K C K P N F Y
Y N G G S P Q* G E A P G V Q* V F A A G A A] A A G V/ I*

SEQ. ID NO 2

GGT GCT GCT TAA GGA GAA GCT AAT GCT AAT TAA CCT TTC GCA GCA AAT AAT GCT
GCT AGA GGT ATA TGT GTA CCA TGC CAA ATA AAC AGA GTA GGC TCT GTT ACC AAT
GCA GGT GAC TTA GCT ACT TTA GCC ACA TAA TGC AGT ACT TAA TGT CCT ACT GGC
ACT GCA CTT GAT GAT GGA GTG ACA GAT GTT TTT GAT AGA TCA GCC GCA TAA TGT
GTT AAA TGC AAA CCT AAC TTT TAC TAT AAT GGT GGT TCT CCT TAA GGT GAA GCT
CCT GGC GTT TAA GTT TTT GCT GCT GGT GCT GCC GCT GCA GGT GTT G

SEQ. ID NO 3

GGA TCC GCT CAG GGA GAA GCT AAT GGT AAT CAG CCT TTC GCA GCA AAT AAT GCT
GCT AGA GGT ATA TGT GTA CCA TGC CAA ATA AAC AGA GTA GGC TCT CGT ACC AAT
GCA GGT GAC TTA GCT ACT TTA GCC ACA CAA TGC AGT ACT CAG TGT CCT ACT GGC
ACT GCA CTT GAT GAT GGA GTG ACA GAT GTT TTT GAT AGA TCA GCC GCA CAG TGT
GTT AAA TGC AAA CCT AAC TTT TAC TAT AAT GGT GGT TCT CCT CAG GGT GAA GCT
CCT GGC GTT CAG GTT TTT GCT GCT GGT GCT GCC GCT GCA CGA ATT C

WHAT IS CLAIMED IS:

1. A vaccine for immunizing fish against ciliated ectoparasitic protozoans comprising an effective amount of a recombinant fusion protein derived from an artificial DNA sequence for immobilization antigen, repeat I, of *Ichthyophthirius multifiliis*.
2. The vaccine according to claim 1 wherein the fusion protein is GST-iAgI.
3. The vaccine according to claim 1 further comprising a substantially inert medium comprising at least one of buffers, adjuvants, immunostimulants or carriers.
4. The vaccine according to claim 1 having the property that, when it is injected into a fish, provides effective protection against white spot disease caused by *Ichthyophthirius multifiliis*.
5. The vaccine according to claim 2, wherein the fusion protein. is produced using E-coli
6. The vaccine according to claim 4, wherein disease is caused by at least one ciliated ectoparasitic protozoan.
7. The vaccine according to claim 6 wherein said ciliated protozoan consists of *Ichthyophthirius multifiliis*.

8. The vaccine according to claim 1 wherein said ciliated protozoans are taxonomically related to *Ichthyophthirius multifiliis*.

9. A method of immunizing fish against ciliated ectoparasitic protozoans comprising treating said fish with an effective amount of a vaccine according to claim 1.

10. The method as claimed in claim 9 wherein said fish are intraperitoneally injected with said vaccine.

11. The method as claimed in claim 9 wherein said fish are fed a medium comprising said vaccine.

12. The method as claimed in claim 9 wherein said fish are dipped in a medium comprising said vaccine.

13. The method as claimed in claim 9 wherein said fish are sprayed with a medium comprising said vaccine.

14. A method of producing a recombinant vaccine against ciliated ectoparasitic protozoans comprising:

a. selecting a region of a amino acid sequence containing a single repeat of *Ichthyophthirius multifiliis* immobilization antigen epitope (i- Ag I);

b. constructing a gene for the selected region from a plurality of synthetic oligonucleotides based on said i-AgI amino acid sequence using universal codons in place of ciliate codons to produce an i-AgI epitope;

c. expressing said i-AgI epitope as a glutathione S transferase (GST) fusion protein in bacteria, *E. coli*; and

d. purifying said GSH-iAgI fusion protein using a glutathione sepharose.

15. The method according to claim 14 wherein said ciliated ectoparasitic protozoans are taxonomically related to *Ichthyophthirius multifiliis*.

16. The method according to claim 14 wherein said ciliated ectoparasitic protozoans comprise *Ichthyophthirius multifiliis*.

17. A method of immunizing fish against ciliated ectoparasitic protozoans comprising treating said fish with an effective amount of a vaccine made according to claim 14.

18. The method as claimed in claim 17 wherein said fish are intraperitoneally injected with said vaccine.

19. The method as claimed in claim 17 wherein said fish are fed a medium comprising said vaccine.

20. The method as claimed in claim 17 wherein said fish are dipped in a medium comprising said vaccine.

21. The method as claimed in claim 7 wherein said fish are sprayed with a medium comprising said vaccine.

22. A synthetic nucleic acid sequence corresponding to SEQ ID No. 3.

23. A synthetic amino acid sequence corresponding to SEQ ID NO. 1

ABSTRACT OF THE DISCLOSURE

A recombinant vaccine that is useful for immunizing fish against ciliated ectoparasitic protozoans that cause white spot disease also known as 'ich'. The disease is caused by *Ichthyophthirius multifiliis* which is a ciliated ectoparasitic protozoan that is an obligate parasite and therefore cannot grow *in vitro*. The instant vaccine replaces chemotherapy and vaccines prepared from *I. Multifiliis* production in live fish. The effective element of the vaccine is a recombinant fusion protein derived from an artificial DNA sequence for immobilization antigen, repeat I, of *Ichthyophthirius multifiliis*. The novel DNA nucleotide and amino acid sequences are disclosed.

Docket No.: 1459-005B

DECLARATION AND POWER OF ATTORNEY

As a below named inventor, I hereby declare that:

My residence, post office and citizenship are as stated below next to my name,

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter claimed and for which a patent is sought on the invention entitled **A RECOMBINANT VACCINE AGAINST FISH INFECTIOUS DISEASES**, the specification of which

☐ is attached hereto ☐ was filed on as Application Serial No. and was amended on (if applicable)

I hereby state that I have reviewed and understand the contents of the above identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose information which is known to me to be material to patentability in accordance with Title 37, Code of Federal Regulations, Section 1.56.

I hereby claim foreign priority benefits under Title 35, United States Code, Section 119(a)-(d) or Section 365(b) of any foreign application(s) for patent or inventor's certificate, or Section 365(a) of any PCT international application which designated at least one country other than the United States, listed below and have also identified below any foreign application for patent or inventor's certificate having a filing date before that of the application on which priority is claimed:

Prior Foreign Application(s):			Priority Claimed	
<u>Number</u>	<u>Country</u>	<u>Day/Month/Year filed</u>	<u>Yes</u>	<u>No</u>
9903188-3	Singapore	28/09/1998	X	

I hereby claim the benefit under 35 USC §119(c) of any United States provisional application(s) listed below.

Prior Provisional Application(s):	
<u>Application Number</u>	<u>Filing Date</u>

I hereby claim the benefit under Title 35, United States Code, Section 120 of any United States application(s), or Section 365(c) of any PCT international application designating the United States, listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States or PCT international application in the manner provided by the first paragraph of Title 35, United States Code, Section 112, I acknowledge the duty to disclose material information as defined in Title 37, Code of Federal Regulations, Section 1.56 which occurred between the filing date of the prior application and the national or PCT international filing date of this application:

Prior U. S. Application(s):		
<u>Serial No.</u>	<u>Filing Date</u>	<u>Status: Patented, Pending, Abandoned</u>

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

I hereby appoint the following attorney(s) and/or agent(s): Allan M. Lowe, Reg. No. 19,641; Israel Gopstein, Reg. No. 27,333; Benjamin J. Hauptman, Reg. No. 29,310; Kenneth M. Berner, Reg. No. 37,093 and Michael G. Gilman, Reg. No. 19,114. all of

LOWE HAUPTMAN GOPSTEIN GILMAN & BERNER, LLP
2034 Eisenhower Avenue, Suite 200
Alexandria, Virginia 22314

with full power of substitution and revocation, to prosecute this application and to transact all business in the Patent and Trademark Office connected therewith, and all future correspondence should be addressed to them.

LOWE HAUPTMAN GOPSTEIN & BERNER, LLP

LOWE HAUPTMAN GOPSTEIN GILMAN & BERNER LLP
DOCKET NUMBER D1459-005B
DECLARATION PAGE 2

Full name of first joint inventor: YOKE MIN SIN

Inventor's signature:

DATE Oct. 1, 1998

Residence: 86 Coronation Road, Singapore 269482

Citizenship: Singaporean

Post Office Address: c/o INTRO, National University of Singapore, 10 Kent Ridge Crescent, Singapore 119260

Full name of second joint inventor: TOONG JIN LAM

Inventor's signature:

DATE Oct. 1, 1998

Residence: 7 Second Avenue, Singapore

Citizenship: Singaporean

Post Office Address: c/o INIRO, National University of Singapore, 10 Kent Ridge Crescent, Singapore 119260

Full name of third inventor: ZHIYUAN GONG

Inventor's signature:

DATE Oct.1, 1998

Residence: 113 Clementi Road #07-01, Singapore

Citizenship: Canadian

Post Office Address: c/o INTR0, National University of Singapore, 10 Kent Ridge Crescent, Singapore 119260

Genotype	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024	2025	2026	2027	2028	2029	2030
1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024	2025	2026	2027	2028	2029	2030	